

Ecologically distinct dinosaurian sister group shows early diversification of Ornithodira

Sterling J. Nesbitt^{1,2,†}, Christian A. Sidor³, Randall B. Irmis^{4,5}, Kenneth D. Angielczyk⁶, Roger M. H. Smith⁷ & Linda A. Tsuji⁸

The early evolutionary history of Ornithodira (avian-line archosaurs) has hitherto been documented by incomplete (*Lagerpeton*¹) or unusually specialized forms (pterosaurs and *Silesaurus*²). Recently, a variety of *Silesaurus*-like taxa have been reported from the Triassic period of both Gondwana and Laurasia, but their relationships to each other and to dinosaurs remain a subject of debate^{3–5}. Here we report on a new avian-line archosaur from the early Middle Triassic (Anisian) of Tanzania. Phylogenetic analysis places *Asilisaurus kongwe* gen. et sp. nov. as an avian-line archosaur and a member of the Silesauridae, which is here considered the sister taxon to Dinosauria. Silesaurids were diverse and had a wide distribution by the Late Triassic, with a novel ornithodiran bauplan including leaf-shaped teeth, a beak-like lower jaw, long, gracile limbs, and a quadrupedal stance. Our analysis suggests that the dentition and diet of silesaurids, ornithischians and sauropodomorphs evolved independently from a plesiomorphic carnivorous form. As the oldest avian-line archosaur, *Asilisaurus* demonstrates the antiquity of both Ornithodira and the dinosaurian lineage. The initial diversification of Archosauria, previously documented by crocodylian-line archosaurs in the Anisian⁶, can now be shown to include a contemporaneous avian-line radiation. The unparalleled taxonomic diversity of the Manda archosaur assemblage indicates that archosaur diversification was well underway by the Middle Triassic or earlier.

By the Middle Triassic, Archosauria had diverged into two lineages: the crocodylian line (Pseudosuchia) and the avian line (Ornithodira, including dinosaurs). During the early evolutionary history of these two clades, pseudosuchians were both numerically abundant and morphologically diverse. In contrast, the avian lineage only became dominant during the Early Jurassic, roughly 35 million years after the origin of Dinosauria^{4,7}.

Triassic deposits have recently begun to yield a diversity of close dinosaur relatives from around the world, whereas records were previously restricted to a single locality from the Middle Triassic of Argentina⁸. New records include lagerpetids, a clade of small forms with long, gracile hindlimbs^{4,9}, and a number of taxa similar to the unusual dinosauriform *Silesaurus*². These groups included contemporaries of dinosaurs, and they persisted well into the Late Triassic, at least until the mid-Norian⁴. *Silesaurus*, from the late Carnian/early Norian of Poland, is unusual among early dinosauromorphs in possessing a beaked lower jaw and a gracile, quadrupedal stance. Here we demonstrate the antiquity, diversity and wide distribution of the clade containing *Silesaurus* and its relatives by introducing a new avian-line archosaur from the Anisian (early Middle Triassic) of Tanzania. The discovery of this well represented taxon demonstrates

that early ornithodiran lineages (pterosaurs, various basal dinosauromorphs, and the lineage leading to Dinosauria) had diverged by the late Anisian.

Archosauria Cope, 1869 (*sensu* Gauthier and Padian, 1985)

Ornithodira Gauthier, 1986 (*sensu* Sereno, 1991)

Dinosauriformes Novas, 1992

Silesauridae clade nov.

Definition: the most inclusive clade for Silesauridae contains *Silesaurus opolensis* Dzik, 2003 but not *Passer domesticus* Linnaeus, 1758, *Triceratops horridus* (Marsh, 1889) and *Alligator mississippiensis* Daudin, 1801.

Diagnosis. Silesauridae differs from all other archosaurs in possessing the following unique combination of character states: rugose ridge on the anterolateral edges of the supraoccipital; notch ventral to femoral head; straight transverse groove on the proximal surface of the femur; and ilium has a straight ventral margin of the acetabulum (see Supplementary Information).

Asilisaurus kongwe gen. et sp. nov.

Etymology. From *asili*, Swahili for ancestor or foundation, and *sauros*, Greek for lizard; *kongwe*, Swahili for ancient.

Holotype. NMT RB9 (National Museum of Tanzania, Dar es Salaam, Tanzania), the anterior portion of a left dentary with associated tooth (Fig. 1a–d).

Paratype. Material found associated with the holotype, but derived from multiple individuals (Fig. 2 and Supplementary Information). Anterior cervical vertebra (NMT RB21), left scapulocoracoid (NMT RB10), sacrum (NMT RB11), proximal portion of an ischium (NMT RB12), ilium (NMT RB13), proximal portion of a pubis (NMT RB14), anterior portion of a skull (NMT RB15), proximal portion of a left humerus (NMT RB16), left astragalus (NMT RB17), right calcaneum (NMT RB18), proximal portion of a left femur (NMT RB19), right tibia (NMT RB20). Additional material from the type locality referable to *Asilisaurus* is under preparation.

Locality and horizon. Most of the specimens derive from a fluvio-lacustrine mudstone–sandstone sequence in the Lifua Member of the Manda Beds¹⁰, near the town of Litumba Ndyosi, Ruhuhu Basin, Tanzania. The type locality preserves the remains of at least fourteen individuals (based on the number of the second sacral vertebra). Cynodonts (*Aleodon brachyrhamphus* and *Scalenodon angustifrons*), dicynodonts (*Sangusaurus* and *Angonisaurus*), archosauromorphs (*Stenaulorhynchus* and pseudosuchians) and stereospondyl amphibians occur in the immediate area^{11–15}. On the basis of comparison with the tetrapod fauna of the subzone C of the *Cynognathus* assemblage zone of

¹Lamont-Doherty Earth Observatory, Columbia University, Palisades, New York 10964, USA. ²Division of Paleontology, American Museum of Natural History, New York, New York 10024, USA. ³Burke Museum and Department of Biology, University of Washington, Seattle, Washington 98195, USA. ⁴Utah Museum of Natural History, 1390 E. Presidents Circle Salt Lake City, Utah 84112-0050, USA. ⁵Department of Geology & Geophysics, University of Utah, Salt Lake City, Utah 84112-0102, USA. ⁶Department of Geology, The Field Museum, 1400 South Lake Shore Drive, Chicago, Illinois 60605, USA. ⁷Karoo Palaeontology, Iziko: South African Museum, Cape Town 8000, South Africa. ⁸Museum für Naturkunde an der Humboldt-Universität zu Berlin, Invalidenstrasse 43, D-10115 Berlin, Germany. [†]Present address: Jackson School of Geosciences, The University of Texas at Austin, Texas 78712, USA.




Figure 1 | New silesaurid from Tanzania, *Asilisaurus kongwe* (NMT RB9). **a**, Anterior portion of the holotypic dentary in medial view. **b**, Anterior portion of the holotypic dentary in occlusal view. **c**, **d**, Scanning electron micrographs of a dentary crown in occlusal (**c**) and lateral (**d**) views. Scale bars: 1 cm (**a**, **b**); 100 μ m (**c**, **d**). bf, bone fragments; ca, carina; f, foramen; fo, fossa; m, matrix; Mg, Meckelian groove; sy, symphysis; t, tooth.

South Africa^{16–18}, the Lifua Member is considered to be late Anisian in age¹⁸.

Diagnosis. *Asilisaurus* differs from all other archosaurs in possessing the following unique combination of character states: anterior portion of the dentary tapers to a sharp point; teeth absent from the anterior portion of the dentary; teeth ankylosed into the alveoli; distinctly convex dorsal margin of the dentary; Meckelian groove positioned at the dorsoventral midpoint of the medial surface of

the dentary; peg-like teeth with extremely small, poorly developed serrations (see Supplementary Information for differential diagnosis using additional characters preserved on the paratypes).

Description. Nearly the entire skeleton of *Asilisaurus* is known from an accumulation of isolated elements and partially articulated individuals at a single locality. Only portions of the skull and manus remain unknown. Most individuals were probably juveniles because the majority of femora are 60–80% of the maximum size recovered. We estimate that the largest individuals of *Asilisaurus* would measure 2–3 m in length, based on femoral length comparisons with *Silesaurus*.

The skull has similar proportions to that of *Silesaurus*². The anterior portion of the dentary (NMT RB9) tapers to a point like that of both *Silesaurus* and *Sacisaurus*¹⁹, but possesses a convex dorsal edge unlike other silesaurids. Small foramina cover the lateral surface of the anterior portion of the dentary, indicating that it was covered by a rhamphotheca. Medially, the prominent Meckelian groove is positioned at the dorsoventral midpoint of the dentary, in contrast to the more ventral position of *Silesaurus*, *Sacisaurus* and Hayden Quarry specimens⁴. A fossa (Fig. 1b) on the occlusal margin separates the tip of the dentary from the tooth row. The closely packed and socketed teeth are ankylosed to the dentary, a synapomorphy uniting *Silesaurus*, *Sacisaurus*, *Asilisaurus* and specimens from the Hayden Quarry⁴. The two preserved teeth are peg-like, but the tip of each crown bears tiny serrations on its carinae.

The anterior cervical vertebrae are significantly longer than both the axis and the posterior cervical vertebrae, similar to *Lewisuchus*²⁰. The rhomboidal cervical vertebrae lack epiphyses, in contrast to the inferred ancestral condition for Dinosauria²¹. Hyposphenepantra articulations are present in the dorsal vertebrae. *Asilisaurus* possesses only the two primordial sacral vertebrae. The sacral ribs are not co-ossified; however, sacral rib one extends onto the body of the last presacral vertebra, a character also present in *Silesaurus*.

The scapula is tall and narrow like that of *Silesaurus*. The glenoid is directed posteroventrally as in dinosaurs, suggesting that the forelimbs were directed underneath the body. The coracoid bears an enlarged biceps tubercle that is separated from the glenoid by a gap. The proximal portion of the humerus is mediolaterally




Figure 2 | Skeletal anatomy of *Asilisaurus kongwe*. **a**, Anterior cervical vertebra (NMT RB21). **b**, Left scapulocoracoid (NMT RB10). **c**, Sacrum (NMT RB11) (reversed). **d**, Proximal portion of the pubis (NMT RB14) (reversed). **e**, Ilium (NMT RB13) (reversed). **f**, Proximal portion of the ischium (NMT RB12) (reversed). **g**, Anterior portion of the skull (NMT RB15) (reversed). **h**, Proximal portion of the left humerus in posterior view (NMT RB16). **i**, Left astragalus in proximal (top) and anterior (bottom) views (NMT RB17). **j**, Right calcaneum in proximal (top) and lateral (bottom) views (NMT RB18). Arrows in **i** and **j** indicate the anterior direction. **k**, Proximal

portion of a left femur in posteromedial view (NMT RB19). **l**, Right tibia in lateral view (NMT RB20). Panels **a–f** are shown in lateral view. Scale bars: 1 cm (**a–l**); 10 cm for the skeleton. Dark grey bones represent missing elements and missing portions of the skeleton based on *Silesaurus*⁵. 4t, fourth trochanter; a., articulates with; ap, anterior process; as, astragalus; c, coracoid; cn, cnemial crest; cr, crest; d, dentary; dia, diapophysis; dp, deltopectoral crest; f, fibula; g, groove; gl, glenoid; il, ilium; is, ischium; la, lacrimal; n, nasal; no, notch; pa, parapophysis; pf, prefrontal; pu, pubis; sar, supra-acetabular rim; sc, scapula; sr, sacral rib; sv, sacral vertebra; ti, tibia; tu, tuber.

expanded like that of *Euparkeria*, and the apex of the deltopectoral crest of the humerus is located at the proximal surface. Although incomplete, the estimated lengths of the humerus and radius indicate that *Asilisaurus* had elongate forelimbs like *Silesaurus*.

The acetabulum of *Asilisaurus* was closed, as with all non-dinosaurian ornithodirans. A distinct crest on the ilium extends dorsally from a well-developed supra-acetabular rim to the pre-acetabular process, as in *Pseudolagosuchus*²², *Silesaurus* and *Sacisaurus*. The postacetabular process of the ilium lacks a brevis shelf. The pubis is anteroventrally directed and terminates as a thick, mediolaterally wide sheet of bone. A groove on the proximal surface of the pubis separates the articular facet with the ilium from the articular facet of the ischium, as in *Silesaurus* and dinosaurs such as *Saturnalia*⁴. The ischial shafts are closely appressed to one another for their entire length and terminate in a posterodorsal expansion.

The femur of *Asilisaurus* possesses a distinct notch ventral to its head. The proximal surface is incised by a straight sulcus similar to that in many basal archosaurs, and like all dinosauriforms, both anteromedial and posterior proximal medial tubera are developed and a distinct facies articularis antitrochanterica is present. Like other basal dinosauriforms⁹, a distinct anterior trochanter is connected to the shaft for the length of the trochanter and bears a poorly developed trochanteric shelf. The crista tibiofibularis is poorly expanded and is separated from the lateral condyle of the femur by a slight groove. The popliteal fossa between the posterior condyles of the femur expands proximally to greater than one-third of the femoral length, a character also present in *Silesaurus*, *Eucoelophysis* and *Sacisaurus*. The tibia bears a well-developed, straight cnemial crest with two equally sized posterior condyles of the proximal portion, similar to other basal dinosauriforms. The distal surface possesses a poorly developed slot for the ascending process of the astragalus.

The astragalus and calcaneum are separate elements, unlike the co-ossified condition in pterosaurs, lagerpetids and theropod dinosaurs. The distinct anterior ascending process of the astragalus is low and poorly developed compared to those of dinosaurs and *Silesaurus*, but nearly identical to the condition in *Pseudolagosuchus*²². A posterior groove is clearly present on the astragalus. The articulation between the astragalus and the calcaneum is much like that of pseudosuchians: the astragalus has a convex surface whereas the calcaneum has a corresponding concave surface. Furthermore, the calcaneum bears a small posterolaterally directed tuber, like that of *Pseudolagosuchus*²² and *Marasuchus*²³. Additionally, the articular facet for the fibula is convex, in contrast to the concave facet of *Silesaurus*³ and Dinosauria.

Our phylogenetic analysis (34 taxa, 290 characters; see Supplementary Information) of *Asilisaurus*, representative pseudosuchians, a diversity of basal ornithodirans, and early dinosaurs recovers *Asilisaurus* in a monophyletic Silesauridae containing *Lewisuchus/Pseudolagosuchus* (see Supplementary Information), *Eucoelophysis*^{3,4,24}, *Sacisaurus*¹⁹ and *Silesaurus*². Silesauridae is supported by four unambiguous synapomorphies (Supplementary Information), whereas the unnamed clade containing *Asilisaurus*, *Eucoelophysis*, *Sacisaurus* and *Silesaurus* is supported by the following synapomorphies: anterior extent of the dentary tapers to a sharp point; teeth absent in the anterior portion of the dentary; maxillary and dentary crowns short and sub-triangular; straight medial articular facet of the proximal portion of the femur; and popliteal fossa of the distal end of the femur extends proximally between one-fourth and one-third the length of the shaft. These synapomorphies allow the placement of more fragmentary specimens from western North America into Silesauridae using unambiguous synapomorphies (Supplementary Information), thus demonstrating that the clade had a Gondwanan distribution during the Middle Triassic and became more widely distributed by the Late Triassic (Fig. 3).

Notably, Silesauridae is well supported as the closest sister taxon of Dinosauria (see Supplementary Information), conclusively demonstrating that *Silesaurus* and *Silesaurus*-like forms are not basal members




Figure 3 | Phylogenetic relationships of *Asilisaurus kongwe* within Archosauria and silesaurid biogeography. Strict consensus of six trees of an analysis with 33 taxa and 290 characters (Supplementary Information). Pseudosuchia, Ornithischia, Sauroptodomorpha, Theropoda, Pterosauroomorpha and Lagerpetidae have been collapsed for clarity. Stars indicate lineages with herbivorous or omnivorous diets. The length of the white bars indicates stratigraphic imprecision. Important synapomorphies for *Asilisaurus* plus *Silesaurus* illustrated on a left femur of *Silesaurus* in posteromedial view (top) and a right dentary of *Silesaurus* in medial view (bottom): A.1, anterior portion of the dentary tapers to a point; A.2, teeth fused to the bone of attachment; A.3, division of the distal condyles of the femur divided for more than one-quarter length of the element; A.4, flat articular facet of the femur. Important synapomorphies for *Sacisaurus* plus *Silesaurus* (B): B.1, Meckelian groove extends through dentary symphysis; B.2, Meckelian groove restricted to the ventral margin of dentary; B.3, teeth with constricted roots. In, Induan; Olen, Olenekian; Rht, Rhaetian. Middle Triassic (squares): 1, *Asilisaurus*; 2, *Lewisuchus/Pseudolagosuchus*. Late Triassic taxa (circles): 3, *Silesaurus*; 4, Eagle Basin specimens; 5, Petrified Forest N.P. taxon; 6, Otis Chalk taxon; 7, *Technosaurus*; 8, *Eucoelophysis*; 9, *Sacisaurus*. Palaeogeographic globe after <http://jan.ucc.nau.edu/~rcb7/globaltext2.html>. See Supplementary Information for further discussion of occurrences.

of the ornithischian dinosaur lineage⁵. All silesaurids lack classical dinosaurian character states such as a laterally open acetabulum, an elongate deltopectoral crest of the humerus, and an extension of the supratermporal fossa onto the posterodorsal surface of the frontal. Thus, features present in *Silesaurus*, such as the ornithischian-like dentition and a theropod-like ankle, represent independent acquisitions (homoplasies) and are not synapomorphies shared with Dinosauria, because these features are not present in the most basal members of Silesauridae (that is, *Asilisaurus* and *Lewisuchus/Pseudolagosuchus*).

Silesaurids possess an unexpected bauplan previously unknown among basal ornithomirans. All known silesaurid taxa represented by adequate material possess an elongate neck, elongate forelimbs and a robust pes. Perhaps the most striking feature of all silesaurids (except *Lewisuchus*) is the presence of teeth with sub-triangular crowns and a constricted root, and dentaries with a beak-like anterior tip, indicating that these taxa were specialized for an omnivorous and/or herbivorous diet^{25,26}. Similar specializations are present among ornithischian, sauropodomorph and some coelurosaurian²⁶ dinosaurs. Because some of the earliest members of these clades lack characters suggestive of a omnivorous and/or herbivorous diet (for example, *Lewisuchus*²⁰ and *Panphagia*²⁷), our analyses suggest that these specializations were independently acquired in at least the silesaurid and sauropodomorph lineages, and possibly in ornithischians as well, and that the shift towards omnivory/herbivory in these three lineages comprised overlapping but distinct suites of morphological characters (Supplementary Information).

The phylogenetic position of *Asilisaurus* within avian-line archosaurs indicates that the radiation of ornithomirans commenced by the early Middle Triassic. Indeed, pterosauriform, lagerpetid, *Marasuchus*, silesaurid and dinosaurian lineages must have all diverged by the late Anisian (Fig. 3). Moreover, the ghost lineage (see Fig. 3) inferred from our phylogeny predicts that taxa closer to Dinosauria than to Silesauridae will be discovered in Middle Triassic strata. Ornithomiran diversification was part of a more inclusive archosaur radiation as evidenced by the diversity of crocodylian-line archosaurs co-occurring in the Manda beds of Tanzania. The Manda archosaur assemblage includes 'rauisuchians'^{13,28}, an *Erpetosuchus*-like taxon²⁹, *Hypselorachis*³⁰, and other undescribed pseudosuchians¹⁵. The minimum of six archosaur lineages represented within the Manda beds is unmatched in similarly aged deposits¹⁸ and foreshadows the taxonomic composition of archosaur-dominated assemblages later in the Triassic. Thus, *Asilisaurus* and other coeval archosaurs from the Manda beds preserve direct evidence that archosaur diversification commenced by the Middle Triassic, but that the group as a whole did not come to dominate terrestrial ecosystems until the Late Triassic.

Received 16 September; accepted 1 December 2009.

- Romer, A. S. The Chañares (Argentina) Triassic reptile fauna. X. Two new but incompletely known long-limbed pseudosuchians. *Breviora* **378**, 1–10 (1971).
- Dzik, J. A beaked herbivorous archosaur with dinosaur affinities from the early Late Triassic of Poland. *J. Vertebr. Paleontol.* **23**, 556–574 (2003).
- Ezcurra, M. D. A review of the systematic position of the dinosauriform archosaur *Eucoelophysis baldwini* Sullivan & Lucas, 1999 from the Upper Triassic of New Mexico, USA. *Geodiversitas* **28**, 649–684 (2006).
- Irmis, R. B. et al. A Late Triassic dinosauriform assemblage from New Mexico and the rise of dinosaurs. *Science* **317**, 358–361 (2007).
- Dzik, J. & Sulej, T. A review of the early Late Triassic Krasiejów biota from Silesia, Poland. *Palaeontologia Polonica* **64**, 1–27 (2007).
- Nesbitt, S. J. *Arizonasaurus* and its implications for archosaur divergence. *Proc. R. Soc. Lond. B* **270** (Suppl.), S234–S237 (2003).
- Brusatte, S. L. et al. Superiority, competition, and opportunism in the evolutionary radiation of dinosaurs. *Science* **321**, 1485–1488 (2008).
- Novas, F. E. Dinosaur monophyly. *J. Vertebr. Paleontol.* **16**, 723–741 (1996).
- Nesbitt, S. J. et al. Hindlimb osteology and distribution of basal dinosauriforms from the Late Triassic of North America. *J. Vertebr. Paleontol.* **29**, 498–516 (2009).
- Wopfner, H. Tectonic and climatic events controlling deposition in Tanzanian Karoo basins. *J. Afr. Earth Sci.* **34**, 167–177 (2002).
- Houghton, S. H. On a collection of Karoo vertebrates from Tanganyika Territory. *Q. J. Geol. Soc. Lond.* **88**, 634–671 (1932).

- von Huene, F. *Stenaulorhynchus*, ein Rhynchosauride der ostafrikanischen Obertrias. *Nova Acta Leopold.* **6**, 7–121 (1938).
- von Huene, F. Ein kleiner Pseudosuchier und ein Saurischier aus den ostafrikanischen Mandaschichten. *Neues Jb. Miner. Geol. Paläont. B* **81**, 61–69 (1939).
- Crompton, A. W. On some Triassic cynodonts from Tanganyika. *Proc. Zool. Soc. Lond.* **125**, 617–669 (1954).
- Sidor, C., Angielczyk, K., Hopson, J., Nesbitt, S. & Smith, R. New vertebrate fossils from the Permo-Triassic Ruhuhu Basin of Tanzania. *J. Vertebr. Paleontol.* **28** (Suppl.), 142A–143A (2008).
- Gay, S. A. & Cruickshank, A. R. I. Biostratigraphy of the Permian tetrapod faunas from the Ruhuhu Valley, Tanzania. *J. Afr. Earth Sci.* **29**, 195–210 (1999).
- Catuneanu, O. et al. The Karoo basins of south-central Africa. *J. Afr. Earth Sci.* **43**, 211–253 (2005).
- Rubidge, B. S. Re-uniting lost continents—fossil reptiles from the ancient Karoo and their wanderlust. *S. Afr. J. Geol.* **108**, 135–172 (2005).
- Ferigolo, J. & Langer, M. C. A Late Triassic dinosauriform from south Brazil and the origin of the ornithischian predatory bone. *Hist. Biol.* **19**, 23–33 (2007).
- Romer, A. S. The Chañares (Argentina) Triassic reptile fauna. XIV. *Lewisuchus admixtus*, gen. et sp. nov., a further thecodont from the Chañares beds. *Breviora* **390**, 1–13 (1972).
- Langer, M. C. & Benton, M. J. Early dinosaurs: a phylogenetic study. *J. Syst. Palaeontology* **4**, 309–358 (2006).
- Arcucci, A. Un nuevo Lagosuchidae (Thecodontia-Pseudosuchia) de la fauna de Los Chañares (edad reptil Chañarense, Triasico Medio), La Rioja, Argentina. *Ameghiniana* **24**, 89–94 (1987).
- Novas, F. E. The tibia and tarsus in Herrerasauridae (Dinosauria, incertae sedis) and the origin and evolution of the dinosaurian tarsus. *J. Paleontol.* **63**, 677–690 (1989).
- Nesbitt, S. J., Irmis, R. B. & Parker, W. G. A critical re-evaluation of the Late Triassic dinosaur taxa of North America. *J. Syst. Palaeontology* **5**, 209–243 (2007).
- Barrett, P. M. Prosauropod dinosaurs and iguanas: speculations on the diets of extinct reptiles. In *Evolution of Herbivory in Terrestrial Vertebrates: Perspectives from the Fossil Record* (ed. Sues, H.-D.) 42–78 (Cambridge Univ. Press, 2000).
- Zanno, L. E., Gillette, D. D., Albright, L. B. & Titus, A. L. A new North American therizinosaurid and the role of herbivory in 'predatory' dinosaur evolution. *Proc. R. Soc. B* **276**, 3505–3511 (2009).
- Martinez, R. N. & Alcober, O. A. A basal sauropodomorph (Dinosauria: Saurischia) from the Ischigualasto Formation (Triassic, Carnian) and the early evolution of Sauropodomorpha. *PLoS One* **4**, e4397 (2009).
- Charig, A. J. *New Triassic archosaurs from Tanganyika including Mandasuchus and Teleocrater*. Abstracts of Dissertations of the Univ. Cambridge 1955–56, 28–29 (1957).
- Walker, A. D. A revision of the Jurassic crocodile *Hallopus*, with remarks on the classification of crocodiles. *Phil. Trans. R. Soc. Lond. B* **257**, 323–372 (1970).
- Butler, R. J. et al. A possible ctenosauriscid archosaur from the Middle Triassic Manda Beds of Tanzania. *J. Vertebr. Paleontol.* **29**, 1022–1031 (2009).

Supplementary Information is linked to the online version of the paper at www.nature.com/nature.

Acknowledgements We thank L. Herzog for preparation of the material and P. Barrett, P. Makovicky, A. Turner and N. Smith for discussions. We acknowledge A. Tibajuka and C. Saanane for their help with permits and fieldwork logistics. Funding was provided by a National Geographic Society Research and Exploration grant (7587-05 to C.A.S.), the Evolving Earth Foundation (to S.J.N.), The Grainger Foundation (to K.D.A.), an NSF Postdoctoral Research Fellowship (DBI-0306158 to K.D.A.), and NSF Graduate Research Fellowships (to S.J.N. and R.B.I.).

Author Contributions S.J.N. and R.B.I. designed the research project; C.A.S. and K.D.A. designed the field project; S.J.N., C.A.S., K.D.A., R.M.H.S. and L.A.T. conducted fieldwork; S.J.N. described the material; R.M.H.S. interpreted the sedimentology and geology; S.J.N. and R.B.I. conducted the phylogenetic analysis; K.D.A. and R.B.I. conducted the comparative phylogenetic analyses; and S.J.N., R.B.I., C.A.S., R.M.H.S., L.A.T. and K.D.A. wrote the manuscript.

Author Information Reprints and permissions information is available at www.nature.com/reprints. The authors declare no competing financial interests. Correspondence and requests for materials should be addressed to S.J.N. (nesbitt@jsg.utexas.edu).